arXiv:0711.0127v1 [nucl-th] 1 Nov 2007

Phenomenological Scaling of Rapidity Dependence for Anisotropic Flows in 25 MeV/nucleon Ca + Ca by Quantum Molecular Dynamics Model

T. Z. Yan,[1, 2] Y. G. Ma,[1, *] X. Z. Cai,[1] D. Q. Fang,[1] G. C. Lu,[1] W. Q. Shen,[1] W. D. Tian,[1] H. W. Wang,[1] and K. Wang[1]
[1]*Shanghai Institute of Applied Physics, Chinese Academy of Sciences, Shanghai 201800, China*
[2]*Graduate School of the Chinese Academy of Sciences, Beijing 100080, China*

(Dated: February 2, 2008)

Anisotropic flows (v_1, v_2, v_3 and v_4) of light fragments up till the mass number 4 as a function of rapidity have been studied for 25 MeV/nucleon ^{40}Ca + ^{40}Ca at large impact parameters by Quantum Molecular Dynamics model. A phenomenological scaling behavior of rapidity dependent flow parameters v_n (n = 1, 2, 3 and 4) has been found as a function of mass number plus a constant term, which may arise from the interplay of collective and random motions. In addition, $v_4/{v_2}^2$ keeps almost independent of rapidity and remains a rough constant of 1/2 for all light fragments.

PACS numbers: 25.75.Ld, 24.10.-i, 21.60.Ka

Collective flow is an important observable in heavy ion collisions and it can bring some information on nuclear equation of state and in-medium nucleon-nucleon cross section [1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15]. In intermediate energy heavy ion collisions, flows can reflect the interplay of collective and random motions. For a thermalized system, the random motions of emitted fragments are dictated by thermal energy, which is independent of mass. In contrast, fragment energy increases linearly with mass if there exists collective motion [16, 17, 18]. Many studies of the properties of the directed flow (v_1) and the elliptic flow (v_2) have been carried out and much interesting physics has been demonstrated on the properties and origin of the collective motion in both nucleonic or partonic levels. Recently, we carried out a Quantum Molecular Dynamics model calculation and found that there is a nucleon-number scaling for the transverse momentum dependent elliptic flow for light particles up to the mass number A = 4 [19, 20, 21]. But there are little studies about higher order flows such as v_3 and v_4 as well as their rapidity dependence. In this work, we shall present rapidity dependence of v_1, v_2, v_3 and v_4 flows for a symmetric collision, namely ^{40}Ca + ^{40}Ca collisions at 25 MeV/nucleon and large impact parameters (b = 4-6 fm) simulated by Quantum Molecular Dynamics model. The scaling behaviors of $v_4/(v_2)^2$ as a function of rapidity is also presented.

Anisotropic flow is defined as the different n-th harmonic coefficient v_n of the Fourier expansion for the particle invariant azimuthal distribution [2]

$$\frac{dN}{d\phi} \propto 1 + 2\sum_{n=1}^{\infty} v_n \cos(n\phi), \tag{1}$$

where ϕ is the azimuthal angle between the transverse momentum of the particle and the reaction plane. Note that the z-axis is defined as the direction along the beam and the impact parameter axis which is labelled as x-axis. Anisotropic flows generally depend on both particle transverse momentum and rapidity, and for a given rapidity the anisotropic flows at transverse momentum p_t ($p_t = \sqrt{p_x^2 + p_y^2}$) can be evaluated according to

$$v_n(p_t) = \langle cos(n\phi)\rangle, \tag{2}$$

where $\langle \cdots \rangle$ denotes average over the azimuthal distribution of particles with transverse momentum p_t. The anisotropic flows v_n can further be expressed in term of single-particle averages:

$$v_1 = \langle cos\phi\rangle = \langle \frac{p_x}{p_t}\rangle, \tag{3}$$

$$v_2 = \langle cos(2\phi)\rangle = \langle \frac{p_x^2 - p_y^2}{p_t^2}\rangle, \tag{4}$$

$$v_3 = \langle cos(3\phi)\rangle = \langle \frac{p_x^3 - 3p_x p_y^2}{p_t^3}\rangle, \tag{5}$$

$$v_4 = \langle cos(4\phi)\rangle = \langle \frac{p_x^4 - 6p_x^2 p_y^2 + p_y^4}{p_t^4}\rangle, \tag{6}$$

where p_x and p_y are, respectively, projections of particle transverse momentum in and perpendicular to the reaction plane.

The intermediate energy heavy-ion collision dynamics is complex since both mean field and nucleon-nucleon collisions play the competition roles. Furthermore, the isospin dependent role should be also incorporated for asymmetric reaction systems. Isospin dependent Quantum Molecular Dynamics model (IDQMD) has been affiliated with isospin degrees of freedom with mean field and nucleon-nucleon collision [22, 23, 24, 25, 26]. The IDQMD model can explicitly represent the many body state of the system and principally contain correlation effects to all orders and all fluctuations, and can describe

---

[*] Corresponding author. Email: ygma@sinap.ac.cn

the time evolution of the colliding system well. When the spatial distance (Δr) is closer than 3.5 fm and the momentum difference (Δp) is smaller than 300 MeV/c between two nucleons, two nucleons can coalesce into a cluster [22]. With this simple coalescence mechanism which has been extensively applied in transport theory, different size clusters can be recognized.

In the model the nuclear mean-field potential is parameterized as

$$U(\rho,\tau_z) = \alpha(\frac{\rho}{\rho_0}) + \beta(\frac{\rho}{\rho_0})^{\gamma} + \frac{1}{2}(1-\tau_z)V_c$$

$$+ C_{sym}\frac{(\rho_n-\rho_p)}{\rho_0}\tau_z + U^{Yuk} \qquad (7)$$

where ρ_0 is the normal nuclear matter density ($0.16 fm^{-3}$), ρ_n, ρ_p and ρ are the neutron, proton and total densities, respectively. τ_z is z-th component of the isospin degree of freedom, which equals 1 or -1 for neutrons or protons, respectively. The coefficients α, β and γ are parameters for nuclear equation of state. C_{sym} is the symmetry energy strength due to the density difference of neutrons and protons in nuclear medium, which is important for asymmetry nuclear matter (here $C_{sym} = 32$ MeV is used), but it is trivial for the symmetric system studied in the present work. V_c is the Coulomb potential and U^{Yuk} is Yukawa (surface) potential. In this work, we take $\alpha = 124$ MeV, $\beta = 70.5$ MeV and $\gamma = 2$ which corresponds to the so-called hard EOS with an incompressibility of $K = 380$ MeV.

About 60000 events have been simulated for the collision system of ^{40}Ca + ^{40}Ca at 25 MeV/nucleon with impact parameter from 4 fm to 6 fm. The physics results were extracted at the time of 200 fm/c when the system has been in the freeze-out stage. Fig. 1(a) shows rapidity dependence of the average in-plane transverse momentum per nucleon $\langle p_x/A\rangle$ around mid-rapidity for light fragments. Here rapidity y is the rapidity of fragment in the center of momentum (c.m.) frame which is normalized to the initial projectile rapidity, i.e., $y = y_{c.m.}/y_{proj}$. It shows that $\langle p_x/A\rangle$ decreases monotonously with the increasing rapidity, and decreases more rapidly for heavier fragments. The negative slope is mainly driven by the attractive mean field in this energy region [11, 27]. Fig. 1(b) shows mass dependence of the so-called sidewards flow parameter (slope value) $d\langle p_x/A\rangle/dy$, which can be easily extracted via linear fits near mid-rapidity in Fig. 1(a). The flow parameter is negative and its absolute value increases with mass apparently, which seems consistent with coalescence-like cluster production mechanism [16].



FIG. 1: (a) The average in-plane transverse momentum per nucleon as a function of the normalized rapidity in the c.m. frame. (b) Mass dependence of $d\langle p_x/A\rangle/dy$. Squares represent for neutrons, circles for protons, triangles for fragments of A=2, diamonds for A=3 and stars for A=4. Error bars show the errors of the fits.



FIG. 2: (a), (b), (c), (d) show rapidity dependence of v_1, v_2, v_3, v_4 for light fragments respectively. Squares represent for neutrons, circles for protons, triangles for fragments of A=2, diamonds for A=3 and stars for A = 4.

In Fig. 2, we show integrated anisotropic flows v_1, v_2, v_3 and v_4 as a function of rapidity y. The trend of the directed flow v_1 shown in Fig. 2(a) is similar to that of $\langle p_x/A\rangle$, i.e., the directed flow decreases monotonously with the increasing rapidity around mid-rapidity. It is positive in target-like rapidity region and negative in projectile-like rapidity region. And it gets to the extremum near $y = \pm 1.5$, and then the absolute value of directed flow goes down with the increasing $\mid y \mid$. It also shows at a given rapidity the absolute magnitude of v_1 is larger for heavier clusters. Fig. 2(b) presents the rapidity dependence of elliptic flow v_2. It shows that the flow is positive and the curve for a given fragment is symmetric with $y = 0$, and descends monotonously when the



FIG. 3: (a), (b), (c) and (d) show rapidity dependence of v_1, v_2, v_3 and v_4 per nucleon for light fragments, respectively. Squares represent for neutrons, circles for protons, triangles for fragments of A=2, diamonds for A=3 and stars for A=4.



FIG. 4: Same as Fig. 3, but for $v_n/C(A)$ (n=1, 2, 3, 4) versus rapidity.



FIG. 5: The ratio of v_4/v_2^2 for neutrons (squares), proton (circles), the fragments of $A = 2$ (triangles), $A = 3$ (diamonds) and $A = 4$ (stars) versus rapidity y.

$| y |$ increases. Again, heavier fragments have larger elliptic flow at the same rapidity. The rapidity dependences of v_1 and v_2 are similar to the results at RHIC energies for charged hadrons [28], but the interaction level of the matter in the two different energy region are obviously different. In this low energy heavy ion collisions, the positive elliptic v_2 essentially stems from the collective rotational motion [11, 27]. Fig. 2(c) and (d) show rapidity dependence of higher order flows of v_3 and v_4, respectively. The trend of v_3 and v_4 are similar to that of v_1 and v_2, respectively, but they show smaller magnitude of flows for a given particle in comparison with v_1 and v_2, respectively. Nevertheless the magnitude of flows of v_3 and v_4 are not too small and hence higher order flows should not be neglected in this energy region.

For testing if the number of nucleon scaling works for rapidity dependent flows, we plot the flows per nucleon as a function of rapidity which is shown in Fig. 3. It seems that the curves for different fragments do not stay together as we imagined except for v_1/A. One reason is that the effect of random motion which is independent of mass may weaken the mass scaling in rapidity space. For details, if we compare the magnitude of collective flows of other fragments with that of proton or neutron, it shows the flow magnitude of fragments with mass number of 2, 3, and 4 is smaller than proton or neutron for v_1 and v_2, but larger than proton or neutron for v_3 and v_4. However, if we use a mass number dependent function $C(A) = \frac{5}{8}(A+\frac{3}{5})$ instead of A for v_1 and v_2, and use $C(A) = 3(A-\frac{2}{3})$ for v_3 and v_4, and plot $v_n/C(A)$ versus y again which is shown in Fig. 4, we can now see the curves for different fragments coincide with each others excellently except for slight deviation in large rapidity region, i.e. in the spectators region. The coefficient $C(A)$ may be qualitatively understood by two parts: the constant part reflects the effect of random motion which is independent of mass, and the part including mass number A reflects the collective motion which increases linearly with mass [16, 29]. From the mass dependent coefficient of $C(A)$, there are two classes: one for v_1 and v_2, and another for v_3 and v_4. The larger coefficient for v_3 and v_4 may indicate that the contribution of collective motion to higher order flows v_3 and v_4 is larger than that to v_1 and v_2.

The RHIC experimental data demonstrated a scaling relationship between 2nd flow v_2 and n-th flow v_n for hadrons, namely $v_n(p_t) \sim v_2^{n/2}(p_t)$ and $v_4/v_2^2 \sim 1.2$ in the STAR data [30]. And we have already shown the scaling of v_4/v_2^2 as a function of p_t in intermediate energy heavy ion collisions [19, 20, 21], i.e., the ratio of v_4/v_2^2 is independent of transverse momentum and shows a constant value of 0.5 for different light fragments. Similarly, we plot p_t-integrated v_4/v_2^2 as a function of rapidity which is shown in Fig. 5. The figure shows that the ratios of v_4/v_2^2 for different fragments up to $A = 4$ roughly keep a constant of 0.5 in the studied rapidity region except for some fluctuations at large rapidities. Therefore the ratio of v_4/v_2^2 is almost independent of transverse momentum and rapidity and its value is around 0.5. In the frame of relativistic fluid dynamics which works in RHIC energies, the ratio value of 0.5 reflects that the collision system reaches to thermal equilibrium and its subsequent evolution follows the laws of ideal fluid dynamics [31]. A coincident value of 0.5 as relativistic fluid dynamics in the present intermediate energy may also indicate some kinds of thermal equilibrium has been also reached.

To summarize, phenomenological behaviors of anisotropic flows as a function of rapidity for light fragments up till mass number 4 have been investigated by the simulation of 25 MeV/nucleon ^{40}Ca + ^{40}Ca collision in peripheral collisions in the framework of the Quantum Molecular Dynamics model. It was shown that v_1 and v_3 of light fragments decrease monotonously with rapidity from positive value near target-like rapidity to negative value near projectile-like rapidity, while v_2 and v_4 are positive and show Gaussian-like shape with a peak around $y_{c.m.} = 0$. When we plot anisotropic flows per nucleon (v_1/A, v_2/A, v_3/A and v_4/A) versus rapidity for light fragments, the curves of different particles do not stay together perfectly except for v_1/A. But when we plot $v_n/C(A)$ (n = 1, 2, 3, 4) versus rapidity where $C(A)$ is a linear function of the mass number A plus an additional constant term, the curves of different particles collapse onto the same curve. That indicates that the fragment flows may arise from the interplay of collective (a term proportional to mass number in $C(A)$) and random thermal motion (a constant term in $C(A)$) of nucleons. $C(A)$ shows a classification between $v_{1,2}$ and $v_{3,4}$, this might reflect that the different role of collective motion on the different anisotropies in momentum space. Additionally, it was found that the ratio of v_4/v_2^2 are almost independent of rapidity and the value is about 0.5, which implies a thermalization scenario of the ideal fluid-like dynamics could be applied even in such low energy heavy ion collisions. Further theoretical investigations and experimental checks are awaiting.

The work was supported partially by the National Basic Research Program of China (973 Program) under Contract No. 2007CB815004, Shanghai Development Foundation from Science and Technology under Grant Numbers 06JC14082 and 06QA14062, the National Natural Science Foundation of China under Grant No 10535010 and 10775167.

[1] Ollitrault J Y 1992 Phys. Rev. D **46** 229.

[2] Voloshin S, Zhang Y 1996 Z. Phys. C **70** 665.

[3] Sorge H 1997 Phys. Lett. B **402** 251; 1997 Phys. Rev. Lett. **78** 2309; 1999 Phys. Rev. Lett. **82** 2048.

[4] Danielewicz P, Lacey R A, Gossiaux P B et al. 1998 Phys. Rev. Lett. **81** 2438.

[5] Zhang B, Gyulassy M, and Ko C M 1999 Phys. Lett. B **455** 45.

[6] Teaney D and Shuryak E V 1999 Phys. Rev. Lett. **83** 4951.

[7] Kolb P F, Sollfrank J, and Heinz U 2000 Phys. Rev. C **62** 054909.

[8] Zheng Y M, Ko C M, Li B A, and Zhang B 1999 Phys. Rev. Lett. **83** 2534.

[9] Li Z Xand Zhang Y X 2006 AIP Conference Proceeding CP865: P. 276, Nuclear Physics Trends: 6th ChinaCJapan Joint Nuclear Physics Symposium, Shanghai, May 16-22, 2006 (Melville, NY: American Institute of Physics), eds Ma Y G and Ozawa A

[10] Persram D and Gale C 2002 Phys. Rev. C **65** 064611.

[11] Ma Y G et al., 1993 Phys. Rev. C **48** R1492; 1993 Z. Phys. A **344** 469; 1995 Phys. Rev. C **51** 1029; 1995 Phys. Rev. C **51** 3256.

[12] Lukasik J et al. (INDRA-ALDAIN Collaboration) 2004 Phys. Lett. B **608** 223.

[13] Lin Z W and Ko C M 2002 Phys. Rev. Lett. **89** 202302; Greco V, Ko C M, Lévai P 2003 Phys. Rev. C **68** 034904.

[14] Chen J H, Ma Y G, Ma G L et al. 2006 Phys. Rev. C **74** 064902.

[15] Abelev B I et al. [STAR Collaboration] 2007 Phys. Rev. Lett. **99** 112301.

[16] Huang M J, Lemmon R C, Daffin F et al. 1996 Phys. Rev. Lett. **77** 3739 .

[17] Partlan M D et al. 1995 Phys. Rev. Lett. **75** 2100.

[18] Jeong S C et al. 1994 Phys. Rev. Lett. **72** 3468.

[19] Yan T Z, Ma Y G et al. 2006 Phys. Lett. B **638** 50.

[20] Ma Y G 2006 AIP Conference Proceeding CP865: P. 284, Nuclear Physics Trends: 6th ChinaCJapan Joint Nuclear Physics Symposium, Shanghai, May 16-22, 2006 (Melville, NY: American Institute of Physics), eds Ma Y G and Ozawa A

[21] Ma Y G, Yan T Z et al. 2007 Nucl. Phys. A **787** 611c.

[22] Aichelin J 1991 Phys. Rep. **202** 233.

[23] Ma Y G and Shen W Q 1995 Phys. Rev. C **51** 710.

[24] Zhang H Y, Shen W Q, Ma Y G et al. 2002 Eur. Phys. J. A **15** 399.

[25] Wei Y B, Ma Y G et al. 2004 Phys. Lett. B **586** 225.

[26] Ma Y G et al. 2006 Phys. Rev. C **73** 014604.

[27] Wilson W K, Lacey R, Ogilvie C A, Westfall G D 1992 Phys. Rev. C **45** 738.

[28] Adams J et al. 2004 Phys. Rev. Lett. **92** 052302; 2005 Phys. Rev. C **72** 014904; 2005 Phys. Rev. Lett. **95** 122301.

[29] Kunde G J et al. 1995 Phys. Rev. Lett. **74** 38.

[30] Adams J et al. 2004 [STAR Collaboration], Phys. Rev. Lett. **92** 062301.

[31] Borghini N and Ollitrault J Y 2006 , Phys. Lett. B **642** 227.